

VAGINAL INFECTION TREATMENTS

Generic	Brand	Form	Dosing Regimen
VULVOVAGINAL CANDIDIASIS			
butoconazole	Gynazole-1	Rx 2% vaginal crm	1 applicatorful once
clotrimazole	—	OTC 1%, 2% vaginal crm	1 applicatorful at bedtime for 3 nights (2% crm) or 7 nights (1% crm)
fluconazole	Diflucan	Rx 150mg oral tab	150mg once
miconazole	Monistat 1 Combination Pack	OTC 1200mg vaginal insert + 2% topical crm	1 insert once at bedtime; apply topical crm twice daily as needed for up to 7 days
	Monistat 3 Cream	OTC 4% vaginal crm	1 applicatorful daily at bedtime for 3 nights
	Monistat 3 Combination Pack	OTC 200mg vaginal supp + 2% topical crm	1 supp, 1 insert, or 1 applicatorful (4% crm) daily at bedtime for 3 nights; apply topical crm twice daily as needed for up to 7 days
	Monistat 3 Cream Combination Pack	OTC 4% vaginal crm + 2% topical crm	
	Monistat 3 Insert Combination Pack	OTC 200mg vaginal insert + 2% topical crm	
	Monistat 7 Cream	OTC 2% vaginal crm	1 applicatorful daily at bedtime for 7 nights
	Monistat 7 Cream Combination Pack	OTC 2% vaginal crm + 2% topical crm	1 applicatorful daily at bedtime for 7 nights; apply topical crm twice daily as needed for up to 7 days
	nystatin	Nystatin Vaginal Tablets	Rx 100,000 Unit vaginal tabs
terconazole	—	Rx 0.8% vaginal crm,	1 applicatorful or supp at bedtime for 3 nights
		80mg vaginal supp	
tioconazole	Monistat 1 Vaginal Ointment	OTC 0.4% vaginal crm	1 applicatorful at bedtime for 7 nights
	Vagistat-1	OTC 6.5% vaginal oint	1 applicatorful once at bedtime
		OTC 6.5% vaginal oint	1 applicatorful once
BACTERIAL VAGINOSIS			
clindamycin	Cleocin Vaginal Cream	Rx 2% vaginal crm	1 applicatorful for 3–7 nights; in 2nd or 3rd trimester of pregnancy use for 7 nights
	Cleocin Vaginal Ovules	Rx 100mg vaginal supp	1 supp for 3 nights
	Clindesse	Rx 2% vaginal crm	1 applicatorful once
metronidazole	Flagyl ER	Rx 750mg oral tab	750mg once daily for 7 days
	MetroGel Vaginal	Rx 0.75% vaginal gel	1 applicatorful once or twice daily for 5 days
	Nuversa	Rx 1.3% vaginal gel	1 applicatorful once daily at bedtime
	Vandazole	Rx 0.75% vaginal gel	1 applicatorful once daily at bedtime for 5 days
sulfanilamide	AVC	Rx 15% vaginal crm	1 applicatorful once or twice daily for 30 days; avoid after 7th month of pregnancy
tinidazole	Tindamax	Rx 500mg oral tab	2g once daily for 2 days or 1g once daily for 5 days

NOTES

All products are for use on consecutive nights unless otherwise noted. Complete relief may take up to 7 days.

Topical creams are to be applied to the external vulval area as needed.

Not an inclusive list of medications, official indications, and/or dosing details.

Please see drug monograph at www.eMPR.com and/or contact company for full drug labeling.

(Rev. 9/2019)